



HAL
open science

Cadmium migration from nib to testa during cacao fermentation is driven by nib acidification

Ruth Vanderschueren, Jasmien Doevenspeck, Florence Helsen, Sandra Mounicou, Jakob Santner, Jan Delcour, Eduardo Chavez, Erik Smolders

► **To cite this version:**

Ruth Vanderschueren, Jasmien Doevenspeck, Florence Helsen, Sandra Mounicou, Jakob Santner, et al.. Cadmium migration from nib to testa during cacao fermentation is driven by nib acidification. LWT - Food Science and Technology, 2022, 157, pp.113077. 10.1016/j.lwt.2022.113077 . hal-03871576

HAL Id: hal-03871576

<https://hal-univ-pau.archives-ouvertes.fr/hal-03871576>

Submitted on 25 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Cadmium migration from nib to testa during cacao fermentation is driven by nib**
2 **acidification**

3

4 Ruth VANDERSCHUEREN ^{a,*}, Jasmien DOEVENSPECK ^a, Florence HELSEN ^a, Sandra
5 MOUNICOU ^b, Jakob SANTNER ^c, Jan A. DELCOUR ^d, Eduardo CHAVEZ ^e, Erik
6 SMOLDERS ^a

7 ^a Division of Soil and Water Management, KU Leuven, Kasteelpark Arenberg 20, 3001
8 Heverlee, Belgium

9 ^b CNRS, Université de Pau et des Pays de l'Adour, E2S UPPA, Institut des sciences analytiques
10 et de physico-chimie pour l'environnement et les matériaux, UMR 5254, 64000 Pau, France

11 ^c University of Natural Resources and Life Sciences, Vienna, Department of Crop Sciences,
12 Institute of Agronomy, Konrad-Lorenz-Straße 24, A-3430 Tulln, Austria

13 ^d Centre for Food and Microbial Technology, KU Leuven, Kasteelpark Arenberg 20, 3001
14 Heverlee, Belgium

15 ^e Escuela Superior Politécnica del Litoral, ESPOL, Facultad de Ciencias de la Vida, Campus
16 Gustavo Galindo Km. 30.5 Vía Perimetral, P.O. Box 09-01-5863, Guayaquil, Ecuador

17

18

19 ABSTRACT

20 Previous work has shown that cacao nib cadmium (Cd) concentrations decrease during
21 fermentation, but only when reaching sufficiently low nib pH. In this work, lab-scale
22 experiments (5 kg units) with lactic and acetic acid amendments were ineffective at reducing the
23 total nib Cd concentration. In contrast, the water-extractable fraction of the nib Cd concentration
24 clearly increased when the pH was decreased. When single pod derived beans were embedded
25 inside a full-scale fermentation box to monitor the effect of the fermentation effect with high
26 precision, nib Cd concentrations decreased by a factor 1.25 (P-value <0.05) after four days of
27 fermentation. Visualisation of the mobile Cd gradient within beans with LA-ICP-MS (using
28 imprints of transversal cuts exposed to a metal binding gel) revealed that fermentation enhances
29 the Cd mobility in the nibs.

30 KEYWORDS

31 Cadmium

32 Cacao

33 Fermentation

34 Laser ablation imaging (LA-ICP-MS)

35 Organic acids

36

37 1. INTRODUCTION

38 In 2019, the European Commission enforced a regulation which sets the maximum allowed
39 cadmium (Cd) concentrations in cacao-derived consumer products (European Commission
40 2014). Similar limits were also approved by the Codex Alimentarius (Codex Alimentarius
41 Commission 2018). These regulations have a large impact on the cacao industry, especially in
42 Latin America where soils naturally contain elevated Cd concentrations. Larger Cd
43 concentrations have been reported in cacao beans and chocolates from Latin America, compared
44 to other geographical origins such as Africa (Bertoldi et al.2016; Vanderschueren et al. 2019). A
45 recent extensive meta-analysis (Vanderschueren et al. 2021) demonstrated the impact of the Cd
46 regulation on cacao farmers in Latin America. More than 50% of compiled bean Cd data
47 exceeded the unofficial industry threshold for EU export (0.60 mg Cd kg⁻¹). The regulation has
48 prompted researchers to study mitigation strategies to lower the Cd concentrations in cacao-
49 derived products. While research thus far focused mostly on the use of soil amendments
50 (Ramtahal et al. 2019) and selection of cultivars for reduced Cd uptake and translocation (Lewis
51 et al. 2018; Engbersen et al. 2019), the potential of postharvest mitigation has also been pointed
52 out (Meter et al. 2019).

53 The effect of fermentation on bean Cd concentrations was recently demonstrated by
54 Vanderschueren et al. (2020), their results indicated that Cd migrates from the nib to the testa
55 and the mucilage during fermentation. This migration decreased the nib Cd concentration by up
56 to a factor 1.3, but only occurred with sufficient nib acidification, i.e. nib pH <5. This migration
57 is counterintuitive since Cd concentrations in the testa are typically higher than in the nib Cd
58 (Ramtahal et al. 2016; Lewis et al. 2018; Vanderschueren et al. 2020), and migration thus occurs
59 against the total Cd concentration gradient. The authors speculated that the concentration

60 gradient of mobile Cd is opposite to the total concentration gradient, especially in more acid,
61 fermented beans.

62 The objective of this study is to better understand the causes of Cd migration during cacao
63 fermentation. More specifically, this study was set up (i) to reveal whether nib acidification is the
64 driving force behind Cd migration during cacao fermentation; and (ii) to identify whether
65 artificial acidification during fermentation using organic acids typically formed during
66 fermentation (acetic and lactic acid) can be used to lower the Cd concentration in the final
67 product. The work consisted of two lab-scale fermentation experiments with different organic
68 acid treatments, and one micro-fermentation experiment established inside a full-scale
69 commercial fermentation unit. The idea behind the lab-scale fermentation experiments is that the
70 effects of acidity on mobile and total nib Cd concentrations can be disentangled from the other
71 effects in the fermentation process, such as heat-induced cell breakdown or the effects of ethanol
72 penetration. A novel method was adopted to map Cd mobility at sub mm resolution. An
73 imprinting technique was used followed by Laser Ablation Inductively Coupled Plasma Mass
74 Spectrometry (LA-ICP-MS) analysis of the imprint. This visualisation method allows to test the
75 hypothesis that the mobile Cd concentration gradient in cacao beans is opposite to the total Cd
76 concentration gradient.

77 2. MATERIALS AND METHODS

78 **2.1. Lab-scale fermentations with organic acid treatments**

79 2.1.1. Experimental setup and sampling

80 Two separate lab-scale (5 kg) fermentation experiments were executed in a greenhouse in
81 Guayaquil (Ecuador) with administration of lactic and acetic acid before, during, or at the end of

82 fermentation (Table 1). Lab-scale fermentations were executed in planter pots with a flat cone
83 shape (small diameter 17.5 cm, large diameter 23.5 cm, height 22.5 cm). Three to five holes in
84 the bottom of each planter pot ensured drainage of fermentation sweatings. Cacao beans
85 (Nacional cultivar) were provided by local farmers in Manabí and Guayas provinces (Ecuador).
86 Prior to setting up each fermentation experiment, 60 kg fresh beans were deposited on a plastic
87 sheet and homogenised manually before dividing them among the planter pots (5 kg each). The
88 planter pots were placed inside Styrofoam boxes to retain heat. The fermentation lasted 5 (lab-
89 scale fermentation A) or 6 days (lab-scale fermentation B). The cacao mass was mixed manually
90 on day 2 in experiment A, and on days 2 and 5 in experiment B. Treatment doses, treatment time
91 and total fermentation time for both experiments are shown in Table 1. To apply the treatments,
92 the cacao beans were spread out in plastic trays and treatment solutions were applied using spray
93 bottles. The application of acids before fermentation and on the second fermentation day
94 logically disturbed the fermentation process and associated temperature increase. Deionised
95 water treatments were included on days 0 and 2 to assess the effect of liquid application on the
96 overall course of fermentation. Those water treatments allowed to disentangle the effect of liquid
97 treatment from the acidification effect. The treated cacao beans were mixed manually to ensure
98 an even distribution of the treatment components and were redeposited in the plastic planter pots.
99 Samples containing 500 g beans were taken after mixing, at the start of fermentation, at the end
100 of fermentation, and before treatment. Samples of the first experiment (A) thus included day 0,
101 day 2, day 4 for the cacao treated that day, and day 5; samples of the second experiment (B)
102 included day 0, day 5 and day 6. The mucilage was removed manually using paper towels, after
103 which the intact beans were oven-dried for 72 hours at 65 °C. After drying, beans were peeled to
104 separate nib and testa, and both fractions were ground using an electric coffee grinder.

105 2.1.2. Water-extractable Cd

106 The Cd mobility was assessed by measuring Cd concentrations in deionised water extracts of the
107 nibs. Ground nib samples were sieved to obtain a more homogenous particle size distribution
108 (800 µm test sieve, VWR International). Sieved material (0.50 g) was incubated with 4.0 mL
109 deionised water in an end-over-end shaker at 20 °C for 5 days. After incubation, samples were
110 centrifuged (15 minutes at 2000 g, Heraeus Multifuge X3R, Thermo Scientific, Waltham, MA,
111 USA). The supernatant was filtered through a 0.45 µm syringe filter and filtered samples were
112 diluted 200 times prior to Inductively Coupled Plasma Mass Spectrometry (ICP-MS) analysis
113 (described below).

114 Additionally, the effect of artificial acidification on Cd mobility was determined using water-
115 extractable Cd as a proxy. To this end, ground nib and testa material from the blank treatments of
116 experiment A (unfermented samples at day 0 and fermented samples at day 5) were used.
117 Duplicate ground materials (0.50 g for nibs and 0.25 g for testa) were incubated in 12.0 mL
118 deionised water, either or not acidified to pH 4.5 with hydrochloric acid (HCl, 37% w/w). The
119 pH of the contact solutions was adjusted prior to incubation and remained unchanged after
120 incubation (<0.1 pH unit difference). Samples were incubated in an end-over-end shaker,
121 centrifuged, and their supernatant was filtered and diluted prior to ICP-MS analysis, as outlined
122 below.

123 2.1.3. Determining the elemental composition

124 Subsamples containing 100 mg dry nib or testa material were acid digested in an open digestion
125 block for 8 h in 3.0 mL Normatom[®] nitric acid (HNO₃ 67–69% w/w, VWR International,
126 Radnor, PA, USA), reaching a maximum temperature of 130 °C. Digests were brought to a
127 volume of 10.0 mL and diluted ten times with Milli-Q water (18.2 MΩ cm⁻¹) prior to elemental

128 analysis with ICP-MS (Agilent 7700x, Agilent Technologies, Santa Clara, CA, USA). Cadmium
129 concentrations were measured by monitoring the ^{111}Cd isotope in helium (He) collision cell
130 mode, using ^{103}Rh as an online internal standard. The limit of quantification (LOQ) for the
131 analysis was $0.001 \text{ mg Cd kg}^{-1} \text{ dw}$. Duplicate samples were included in the digestions at intervals
132 of ten samples and the coefficient of variation (CV) of these duplicates ranged from 0.4 to 13%
133 (average CV 3.5%). Blank samples (in quadruplicate) and certified reference NIST 2384 baking
134 chocolate (in triplicate, certified concentration $0.073 \pm 0.008 \text{ mg Cd kg}^{-1}$) were included in all
135 digestions and treated identical to the cacao samples. Recoveries of the certified reference
136 material ranged between 87 and 100%. In addition to Cd, several other elements were also
137 analysed (Al, P, K, Ca, Cr, Mn, Co, Ni, Cu, Zn, As, Mo and Pb). Specific parameters regarding
138 the ICP-MS analysis for those elements and a discussion on the quality control can be found in
139 the Supplementary Information (SI).

140 2.1.4. Temperature and pH

141 The temperature (T) in the centre of the fermentation boxes was measured daily using a digital
142 thermometer (VWR International). The pH was determined only at sampling times to avoid
143 excessive disturbance of the fermentation process. The pH of the mucilage, not including the
144 testa, was determined in suspension after shaking 25 g beans with mucilage in 100 mL deionised
145 water on an orbital shaker (KS 130 orbital shaker, IKA laboratory equipment, Staufen, Germany)
146 for 5 minutes at 320 rpm. The nib pH was determined by shaking 5.0 g dried ground material in
147 20 mL deionised water on an orbital shaker for 5 min and measuring the pH in the supernatant
148 after centrifugation for 25 minutes at 1800 g (MTL-5MS Tabletop Low Speed Centrifuge,
149 Micronlab, Shandong, China).

150 2.1.5.

151

152

153 **2.2. Micro-fermentation**

154 2.2.1. Cacao material and setup

155 The micro-fermentation experiment was carried out at a plantation in Guayas province
156 (Ecuador), using Nacional cultivar cacao. Fifteen ripe cacao pods were opened using a machete.

157 Half of the beans from each pod were immediately sampled and processed without fermentation.

158 The other half of the beans from each pod were collected in mesh bags with drawstring closing

159 and placed in the centre of a wooden fermentation box (82 x 65 x 65 cm) filled with Nacional

160 cacao from the same field (225 kg). Fermentation lasted four days and the mass was mixed

161 manually on the second day. The mesh sample bags were kept in the centre of the box

162 throughout the process and were sampled when fermentation was complete (fermented samples)

163 (SI, Figure S 1).

164 Unfermented and fermented beans were cleaned with paper towels to remove the mucilage,

165 oven-dried for 72h at 65 °C, and peeled to separate nib and testa. Due to the small sample size,

166 testa and nibs of three individual beans per sample were digested separately and their elemental

167 composition was analysed with ICP-MS. The temperature in the centre of the fermentation box

168 was measured daily using a digital thermometer (Custom, CT-422WR, Hitchcock, TX, USA).

169 Nib and mucilage pH were determined before and after fermentation in the bulk cacao

170 surrounding the micro-fermentation samples, as described above.

171 2.2.2. Visualisation of Cd mobility using LA-ICP-MS

172 A novel visualisation method was developed to map mobile Cd in cacao beans at mm resolution,
173 using the principles of the Diffusive Gradient in Thin film technique (DGT) developed by
174 Santner et al. (2015). In brief, a transversely cut bean was exposed to a gel that contains a metal
175 ion selective binding agent. The mobile metal ions migrate from the source to the gel. After a
176 given deployment time, the gel is removed, dried, and its surface composition is scanned with
177 LA-ICP-MS.

178 Metal binding gels containing micro-milled Chelex-100 resin were prepared as described by
179 Zhou et al. (2018), with a resin-to-gel ratio of 4.5% (w/w). Due to the small size of the resin
180 beads (<10 µm), this gel can be used for chemical analysis with high spatial resolution. Mobile
181 Cd was visualised for 5 of the 15 micro-fermentation pods and included two unfermented and
182 two fermented beans from each pod. Immediately after sample collection (either before or after
183 fermentation, not dried), beans were transversely cut in half using a clean razor blade and placed
184 on the gel overnight. A Nuclepore[®] membrane (0.2 µm pore size, Whatman Maidstone, UK) was
185 placed in between the beans and the gel as a barrier to avoid particulate transfer. The micro
186 Chelex gel acts as a zero sink for cationic trace metals. It binds mobile Cd ions present in the
187 contact surface with the bean and thereby creates a print of the mobile Cd in the beans. Imprinted
188 Chelex gels were stored in plastic bags with several drops of ultrapure water to keep the gels
189 hydrated. Gels were dried on 0.45 µm Millipore membrane with Whatman chromatography
190 paper as a support (3 mm CHR), in a slab gel dryer (DrygelSr, Hoefer, Holliston, MA, USA) for
191 at least 48 hours. During drying, gels were covered with clean polyethylene plastic to protect the
192 reactive layer. Dried gels were fixed on glass microscopy slides using double sided tape and
193 submitted to LA-ICP-MS analysis.

194 For elemental analysis, a quadrupole 7700cs ICP-MS (Agilent Technologies) mounted with
195 platinum cones was used. The sensitivity and operational conditions of the ICP-MS (stability,
196 background and mass calibration) were initially checked using a 1.0 $\mu\text{g L}^{-1}$ Y, Tl, Li, Ba and Ce
197 tuning solution. The ICP-MS was then coupled to a 213-nm laser ablation system equipped with
198 a TV2 cell (NWR213, ESI, Fremont, CA, USA) and coupling was optimised using a NIST 612
199 glass by monitoring ^{238}U and ^{232}Th for maximal sensitivity, and a U to Th ratio as close as
200 possible to the unit (NIST 612 certified concentrations $37.79 \pm 0.08 \text{ mg Th kg}^{-1}$ and 37.38 ± 0.08
201 mg U kg^{-1}). Visualisation of Ca, Cd, K, Ni and Zn fixed on micro Chelex gel was performed by
202 running several ablation line scans with a 20 Hz laser shot repetition rate, fluency maintained
203 between 4.1 and 4.5 J cm^{-2} , a laser beam of 50 μm^2 and a scan speed of 50 $\mu\text{m s}^{-1}$. Ablated
204 material was transported with 800 mL min^{-1} He and mixed with Ar gas before the ICP torch
205 inlet. The ICP-MS was used in He mode, allowing monitoring of ^{111}Cd (0.2 s), ^{114}Cd (0.2 s), ^{39}K
206 (0.005 s), ^{44}Ca (0.15 s), ^{60}Ni (0.1 s), and ^{64}Zn (0.1 s as integration time).

207 Several sets of two or three ablation lines (distance between lines 100 μm) were run on each
208 sample (SI Figure S 6). Signal intensities were averaged for each set of ablation lines. To account
209 for the differences in elemental composition between samples and elements, signal intensities
210 measured for each element (counts per second, cps) were divided by the elemental concentration
211 determined in each sample by acid digestion and ICP-MS analysis as described above. Several
212 gels were lost prior to LA-ICP-MS imaging as they were either compromised during storage and
213 transportation or broken during gel drying. In addition, gels that displayed disproportionately low
214 K intensities, i.e. factor 10 lower than the other samples, were excluded from the discussion
215 below. Even though the affinity of the Chelex resin is lower for K compared to the other
216 elements evaluated here (Cd, Ca, Ni and Zn), K concentrations in the nibs are high and K is

217 readily water-extractable (see results below). Therefore, low K intensities were considered as an
218 indication for poor contact between gel and bean, or as an indication that the gel was erratically
219 used upside down, as these gels are typically asymmetric because Chelex resin beads settle to the
220 bottom during gel casting.

221 3. RESULTS AND DISCUSSION

222 **3.1. Lab-scale fermentations with organic acid treatments**

223 3.1.1. Effect on fermentation parameters

224 Both water and organic acid treatments affected the fermentation temperature. It dropped to
225 ambient temperature (i.e., 30 °C or below) within 24 hours after treatment in both experiments
226 (Figure 1 and SI, Figure S 2). Pre-fermentation treatment (day 0) delayed heat production in the
227 fermenting mass. In fermentation A, this effect was stronger for the acid compared to the
228 corresponding water treatment (SI, Figure S 2). The temperature profiles of fermentation A
229 suggest that the effect of treatment on temperature was smallest when applying the acids after
230 fermentation, as was performed in fermentation B.

231 The mucilage pH was strongly affected by the treatments as this tissue was in direct contact with
232 the solutions. The final mucilage pH was significantly lower in acid treated cacao compared to
233 control and water treatments in both experiments [Tukey's Honestly Significant Difference
234 (HSD) test, P-value < 0.05, Table 2]. Water treatment on day 0 increased the mucilage pH from
235 4.0 before fermentation to 6.9 on day 5, indicating suboptimal fermentation. Water treatment
236 prior to fermentation likely delayed and/or impeded the outgrowth of yeasts in the first
237 fermentation phase, which reduced the ethanol production. Ethanol is the main substrate for the
238 acetic acid bacteria in the third and final fermentation phase. During this phase, the mucilage pH

239 decreases as these bacteria convert ethanol to acetic acid, and the fermentation temperature
240 increases to >45 °C because this process is strongly exothermic. Thus, reduced production of
241 ethanol may explain the higher final mucilage pH and lower maximum temperature observed in
242 the pre-fermentation water treatment.

243 At the end of experiment A, the nib pH was significantly lower in the control, and in the acetic
244 acid day 2 and day 4 treatments compared to acetic acid day 0 and water treatments (Tukey's
245 HSD test, P-value < 0.05 , Table 2). The acetic acid treatments in experiment A thus did not
246 lower the nib pH below that of the control, and results of the water treatments indicated that
247 liquid application hindered nib acidification with fermentation. This again suggests reduced
248 availability of ethanol substrate for the acetic acid bacteria, as nib acidification mostly results
249 from penetration of acetic acid into the nibs. Acetic acid treatment before fermentation (day 0)
250 immediately decreased the nib pH, but this effect disappeared with time [significant decrease
251 from pH 6.5 ± 0.1 (standard deviation, stdev) to 5.9 ± 0.1 within one hour after treatment,
252 Student's t-test, P-value < 0.05 , results not shown). Despite this immediate penetration of the
253 acetic acid into the nibs, pre-fermentation treatments yielded higher final nib pH values
254 compared to control treatments, which was likely related to the effects of the liquid application
255 on the overall fermentation (i.e. less acetic acid production from fermentation activity).

256 Much larger acid doses were applied in experiment B compared to A (35 and 53 g kg⁻¹ cacao
257 compared to 13 and 15 g kg⁻¹ cacao, Table 1). These treatments were applied only after
258 fermentation and resulted in significantly lower final nib pH values (on average 4.4 for lactic
259 acid and 4.5 for acetic acid treatment) compared to control or water treatments (average nib pH
260 5.2 , Tukey's HSD test, P-value < 0.05 , Table 2). A detailed overview of the changes in nib pH
261 with fermentation time can be found in SI Figure S 3. Within the first hour after application, the

262 average nib pH decreased significantly from 5.3 ± 0.2 (stdev) to 4.6 ± 0.1 with lactic acid and
263 from 5.3 ± 0.1 to 4.7 ± 0.1 with acetic acid application (Students t-test, P-value <0.05).
264 Penetration of both acids into the nib thus occurs quickly and does not require fermentation, as
265 equally fast acid penetration was observed in the pre-fermentation treatment of experiment A. It
266 should be noted that, while both lactic and acetic acid treatments in experiment B resulted in a
267 lower nib pH compared to the blank and water treatments, final nib pH values were still within
268 the range for fermented cacao reported in literature, i.e. pH 4.0–5.5 (Schwan and Wheals 2004;
269 Belitz et al. 2009; Papalexandratou et al. 2011; De Vuyst and Weckx 2016). The differences in
270 final nib pH between acid treated and untreated cacao (control or water) in experiment B thus
271 may have been related to less extensive fermentation in these small planter pots.

272 3.1.2. Effect on tissue Cd concentrations

273 A detailed overview of nib and testa Cd concentrations as a function of fermentation time can be
274 found in SI Figures S 3 and S 4, a summary is given in Table 2. Nib Cd concentrations were not
275 significantly affected by fermentation (Tukey's HSD test, P-value ≥ 0.05), except for the
276 replicates treated with acetic acid on day 2 in experiment A, where nib Cd decreased from $2.3 \pm$
277 $0.3 \text{ mg Cd kg}^{-1}$ before fermentation to $2.0 \pm 0.1 \text{ mg kg}^{-1}$ after fermentation (day 5) (SI, Figures S
278 3 and S 4). In both fermentation experiments, no significant differences could be observed in
279 final nib Cd concentrations among the different treatments (Tukey's HSD test, P-value ≥ 0.05 ,
280 Table 2). However, final testa Cd concentrations did differ significantly among treatments in
281 both experiments. Because the testa represents a much smaller weight fraction than the nib, the
282 total Cd stock in the testa is also smaller. For example, considering the weight fractions of nib
283 and testa from which mucilage had been removed [average weight fractions nib 0.93 and testa
284 0.07 (Vanderschueren et al. 2020)], a cacao bean from the control in experiment A with a total

285 weight of 1 g (nib and testa) would have a total stock of 1.8 μg Cd in the nib but only 0.26 μg Cd
286 in the testa. This explains why outward Cd migration from nib to testa can result in a significant
287 increase in the testa Cd concentration but not in a detectable decrease in the nib Cd
288 concentration. However, patterns were not consistent as testa Cd concentrations increased in
289 some treatments but decreased in others (SI, Figures S 3 and S 4). The inconsistency in the effect
290 of fermentation on testa Cd concentrations may have been related to further migration of Cd
291 toward the mucilage. This hypothesis could not be verified as the mucilage was not collected for
292 elemental analysis in this study.

293 While both lactic and acetic acid treatments in experiment B resulted in a similar final nib pH of
294 4.5, testa Cd increased more extensively with acetic acid compared to lactic acid treatment
295 (Table 2). Both acids thus penetrated and acidified the cacao nib, but testa Cd results suggested
296 that only acetic acid treatment resulted in relevant Cd mobilisation. However, this hypothesis
297 was not confirmed by the water extractions, as the water-extractable Cd fractions in the nibs of
298 both lactic and acetic acid treated cacao were similar at the end of fermentation and did not differ
299 from those in blank or water treatments. In experiment A, the total nib and testa Cd
300 concentrations decreased and increased significantly with decreasing nib pH (Pearson correlation
301 test, P-value < 0.05, Figure 2). In experiment B, no significant correlations could be found. Still,
302 a trend of increasing testa Cd concentrations with decreasing nib pH could be discerned, with the
303 acetic acid treated samples standing out in the low pH/high testa Cd area of the graph (Figure
304 2.B, top left). This relation between tissue Cd and nib pH suggests a pH dependent mobilisation
305 of Cd in the nib tissue. Indeed, the water-extractable nib Cd fraction increased significantly with
306 decreasing nib pH in both experiments (Pearson correlation test, P-value < 0.05). The effects of

307 different fermentation treatments on tissue elemental concentrations and water-extractable nib
308 fractions of Ca, Cu, K, Mn, Ni, P and Zn are reported in SI Table S 1 and Figure S 6.

309 The water extraction of unfermented bean tissues, that were artificially acidified to pH 4.5,
310 indicated an increase of water-extractable nib Cd at lower pH (Figure 3), again suggesting the
311 importance of acidification for Cd migration. Similar effects of acidification were observed for
312 e.g. Mn and Zn (SI, Figure S 7). Water-extractable Cd fractions were much lower in the testa
313 compared to the nib. The increased water-extractability of Cd in the fermented testa may
314 represent the Cd that migrated from nib to testa during fermentation, while the Cd present in this
315 tissue prior to fermentation was less water-extractable (Figure 3).

316 **3.2. Micro-fermentation**

317 3.2.1. Effect on tissue Cd concentrations

318 The temperature in the centre of the box increased from ambient temperature (26 °C) to a
319 maximum of 50 °C on the third day, after which it decreased to 46 °C by day 4. The mucilage
320 pH increased from 3.9 (day 0) to 4.4 (day 4), while the nib pH decreased from 6.3 (day 0) to 4.8
321 (day 4). Even though all pods were collected from a single field, both nib and testa Cd
322 concentrations varied by more than a factor 3 among the pods (CV 37% for nibs and 31% for
323 testa), while variations among beans within a pod were relatively small (average CV 12% for
324 nibs and 10% for testa) (Figure 4). Similar large variations in bean Cd within a single field were
325 found previously by Argüello et al. (2019). They reported an average within-field CV of 39% in
326 a nationwide study in Ecuador. Pairwise analysis of single pod micro-fermentations revealed an
327 effect of fermentation on both nib and testa Cd concentrations. A pairwise t-test indicated a
328 significant decrease in the nib Cd concentration with fermentation by $-0.30 \text{ mg Cd kg}^{-1}$, and an
329 increase in the testa Cd concentration by $+0.75 \text{ mg Cd kg}^{-1}$ (P-value <0.05). In contrast, unpaired

330 analysis of the ensemble of data did not detect these changes (SI Table S 2). The nib Cd
331 concentration decreased on average by a factor 1.25, which is similar to the factor 1.3 reported
332 previously (Vanderschueren et al. 2020). Significant decreases in nib elemental concentrations
333 were also observed for Cu (factor 1.3), K (factor 1.3), Ni (factor 1.3) and P (factor 1.1), while
334 testa elemental concentrations increased for Cu (factor 2.9), K (factor 2.5), Mn (factor 2.6), Ni
335 (factor 2.2), P (factor 5.5) and Zn (factor 2.1) (SI Table S 2).

336 Considering the weight fractions of nib and testa from which mucilage had been removed
337 [average weight fractions nib 0.93 and testa 0.07 (Vanderschueren et al. 2020)], the average mass
338 of Cd lost from the nibs ($0.28 \pm 0.20 \text{ mg Cd kg}^{-1} \text{ bean}$) was significantly larger than the mass of Cd
339 gained in the testa ($0.05 \pm 0.03 \text{ mg Cd kg}^{-1} \text{ bean}$). Similar discrepancies were found in the mass
340 balances of Cu, K, Ni and P, i.e. the decrease in mass of these elements in the nib was
341 significantly larger than the mass gained in the testa. This discrepancy in mass balances is likely
342 related to leaching of elements into the mucilage (the liquid phase surrounding the beans).
343 Indeed, Cd, Cu, K, Ni and P concentrations in the mucilage (among other elements) have been
344 reported to increase with fermentation by a factor 2–8 depending on the element and on the
345 fermentation conditions. It was also reported that fermentation reduced the total bean Cd level by
346 15%, likely due to leaching of Cd in the mucilage sweatings (Vanderschueren et al. 2020).

347 3.2.2. Visualisation of elemental mobility

348 While duplicate mobile elemental prints were originally generated before and after fermentation
349 for five micro-fermentation pods, only eight samples are discussed here because several samples
350 had to be excluded, as explained above. The contact area between the beans and the Chelex resin
351 gels could be clearly discerned on dried gels, as the samples stained the gel (Figure 5 and SI
352 Figure S 8). This imprint allowed targeting the contact area during analysis. Visualisation of the

353 elemental imprints on the Chelex resin gels by LA-ICP-MS indicated that mobility of Cd, Ni and
354 Zn increased with fermentation, with largest mobile element signals located in the centre of the
355 nib imprints (Figure 5). Mobile K also increased while mobile Ca decreased with fermentation
356 (SI Figure S 9). For Cd and Zn, the observations are in accordance with the increased water
357 extractability of these elements in fermented nibs (Figure 3 and SI Figure S 6). The water
358 extractability of Ni in nibs was not significantly affected by fermentation, more than 70% of nib
359 Ni was water-extractable regardless of the nib pH (SI Figure S 6). However, signal intensities for
360 Ni on the Chelex gels were clearly larger for fermented than for unfermented samples, and nib
361 Ni concentrations were positively correlated to nib pH in both fermentation experiments (SI
362 Table S 2 and Figure S 6). The maps of the mobile element prints obtained here (signal larger for
363 nib than for testa) were markedly different from the LA-ICP-MS images of total element
364 composition in unfermented cacao beans obtained previously, where the total Cd signal was
365 larger for the testa than for the nib (Vanderschueren et al. 2020). This suggests that fermentation
366 induces an increase in Cd mobility in the nibs but not in the testa, resulting in a mobile
367 concentration gradient opposite to the total Cd concentration gradient. The lack of element signal
368 peaks at the edges of the Chelex imprints was in accordance with the lower water-extractable Cd
369 fraction found in the testa compared to the nib. Imaging with LA-ICP-MS using imprints of
370 cacao beans on a metal chelating gel was used to successfully visualise element mobility in
371 unfermented and fermented cacao beans. As such, the visualisation method was used here to
372 better understand the underlying mechanisms of Cd migration during cacao fermentation, while
373 bulk chemical analysis using acid digestion and ICP-MS (see above) allows to quantify the
374 extent of migration occurring. This novel imaging technique can be employed in the future for
375 the visualisation of element mobility in a variety of sample matrices and may be strengthened by

376 including analysis of chelating gel standards with predetermined Cd concentrations which would
377 allow (semi-)quantitative analysis of the images.

378 **3.3. Impact of the experimental setup**

379 The results of the micro-fermentation experiment revealed a factor 1.25 decrease in nib Cd
380 (Figure 4), but no such effect was found in the lab-scale experiments (Table 2). The lack of
381 effect in the lab-scale experiments was likely not related to insufficient acidification. The final
382 nib pH for the control in lab-scale fermentation A (pH 4.6) was even lower than that reached in
383 the micro-fermentation (pH 4.8). The difference in effects between setups may be related to
384 differences in temperature. While small lab-scale fermentations of 5 kg cacao have practical
385 advantages for experimental work, the small fermentation volume offers less heat retention
386 compared to micro-fermentation experiments performed in a box of >200 kg cacao. The
387 temperature in the centre of the lab-scale vessels reached 45 °C in the controls of both
388 experiments (A and B, Figure 1 and SI Figure S 2), which was on the lower end of maximum
389 temperatures typically reported in literature (Lima et al. 2011) and lower than the temperature
390 reached in the micro-fermentation (50 °C). The lab-scale temperature profiles showed a delay in
391 initial temperature rise, especially in experiment B where the temperature only started to increase
392 after two days (Figure 1). Jespersen et al. (2005) reported that yeast outgrowth in the outer layers
393 of the fermenting cacao mass is delayed in comparison to that in the centre of the mass. They
394 found that the maximum yeast cell count was reached after 24 hours in the centre, while it was
395 only reached after 72 h in the outer layers. The delay in fermentation activity observed in the lab-
396 scale fermentations, indicated by the lag phase in the temperature profile, may be related to the
397 small size and large surface-to-volume ratio of the lab-scale vessels. Fermentation temperature
398 may thus explain the lack of decrease in nib Cd concentrations in the lab-scale experiments.

399 Sufficient heat may be required to allow acidification-driven mobilisation of Cd in cacao nibs
400 during fermentation. Additional research under controlled temperature conditions is required to
401 verify or reject this hypothesis.

402 The differences in detected effects between both setups may also be related to the large variation
403 in bean Cd. Results of the micro-fermentation experiment demonstrated how large inter-pod
404 variability (up to a factor 3) can mask the effect of fermentation on bean Cd concentrations
405 (Figure 4). The coefficient of variation of the nib Cd concentration between all samples prior to
406 treatment in lab-scale fermentation B was large (CV 13%). A power test was performed to assess
407 the number of replicates in the lab-scale experiments, using the online tool developed by Rollin
408 Brant [University of British Columbia (<https://www.stat.ubc.ca/~rollin/stats/ssize/n2.html>)]. To
409 account for differences in the initial bean Cd concentrations, the CV was used instead of the
410 standard deviation. Based on that analysis, at least four replicates are required to detect a factor
411 1.3 effect with statistical significance (CV = 13%, $\alpha = 5\%$ and power = 80%). However, the
412 power test also indicated that two replicates per treatment were sufficient to detect the effect in
413 lab-scale fermentation A (CV between all nib samples prior to treatment was only 6%).

414 **3.4. Organic acid fermentation treatments as a mitigation strategy for Cd in cacao**

415 The micro-fermentation experiment indicated that nib acidification is the driver for nib Cd
416 mobilisation, and thus also for reducing the final nib Cd concentration. However, organic acid
417 application in lab-scale fermentations did not reduce the nib Cd concentration, probably because
418 of the lower temperatures compared to commercial fermentations. Organic acid treatments
419 reduced the nib pH, but only when applying large doses, and the final nib pH was still within the
420 range of commercial values (pH 4–5). Thus, nib Cd does decrease with decreasing nib pH but
421 organic acid treatments during fermentation may not be the best option to acidify the nibs.

422 Optimisation of fermentation conditions can likely yield final nib pH values similar to those
423 resulting from organic acid treatments. For example, acetic acid concentrations in mucilage are
424 reported to be higher when turning is performed (Camu et al. 2008a). The impact of such
425 fermentation practices on the flavour should be taken into account, as excessive acetic acid has
426 been related to less pronounced flavour in the final product (Camu et al. 2008b).

427 4. CONCLUSION

428 The mobility of Cd in cacao nibs increases during fermentation, as indicated by enhanced water
429 extractions and enhanced diffusible Cd identified by LA-ICP-MS imaging. Water-extractable nib
430 Cd was negatively correlated to nib pH and increased through artificial acidification, indicating
431 nib pH as the driving force behind nib Cd mobilisation. Migration of Cd from the nibs to the
432 testa during fermentation is likely related to increased Cd mobility in the nibs due to
433 acidification. Organic acid amendments before or during fermentation were ineffective in
434 reducing the nib pH due to their negative impact on fermentation (lower fermentation
435 temperatures and excessively high mucilage pH). Application of acetic or lactic acid as a post-
436 fermentation treatment lowered the nib pH compared to blank and water treatments, but only
437 when applying large doses. Although no significant reductions in nib Cd concentrations could be
438 observed with acid treatment, the reduced nib pH did increase testa Cd concentrations and nib Cd
439 mobility. Micro-fermentations with single pod-derived beans inside a commercial fermentation
440 box indicated a factor 1.25 reduction in nib Cd. Optimisation of fermentation parameters for
441 acetic acid production can likely offer similar nib acidification compared to acid treatments,
442 without the added cost. However, a trade-off needs to be made between nib Cd reduction and
443 flavour quality as excessive nib acidity is considered an off-flavour.

444 ACKNOWLEDGEMENTS

445 RV is the recipient of an FWO-SB fellowship (FWO Vlaanderen). JAD receives Methusalem
446 research excellence financing at KU Leuven. Aquitaine Region and Feder are acknowledged for
447 the funding of the laser ablation system from the Aquitrace Project (convention number
448 20131206001-13010973). The authors thank the Ecuadorian farmers who supplied material and
449 allowed us to use their infrastructure. We also thank Dr. David Arguëllo and Jesse Dekeyrel for
450 assistance with the lab work.

451

452 REFERENCES

453 Argüello D, Chavez E, Lauryssen F, Vanderschueren R, Smolders E, Montalvo D. 2019. Soil
454 properties and agronomic factors affecting cadmium concentrations in cacao beans: A
455 nationwide survey in Ecuador. *Sci Total Environ.* 649:120–127.
456 doi:10.1016/j.scitotenv.2018.08.292.

457 Belitz HD, Grosch W, Schieberle P. 2009. Coffee, tea, cocoa. In: *Food Chemistry*. Berlin,
458 Heidelberg: Springer Berlin Heidelberg. p. 938–970. doi:10.1007/978-3-540-69934-7.

459 Bertoldi D, Barbero A, Camin F, Caligiani A, Larcher R. 2016. Multielemental fingerprinting
460 and geographic traceability of *Theobroma cacao* beans and cocoa products. *Food Control*.
461 65:46–53. doi:10.1016/j.foodcont.2016.01.013.

462 Camu N, González Á, De Winter T, Van Schoor A, De Bruyne K, Vandamme P, Takrama JS,
463 Addo SK, De Vuyst L. 2008a. Influence of turning and environmental contamination on the
464 dynamics of populations of lactic acid and acetic acid bacteria involved in spontaneous cocoa
465 bean heap fermentation in Ghana. *Appl Environ Microbiol.* 74(1):86–98.
466 doi:10.1128/AEM.01512-07.

467 Camu N, De Winter T, Addo SK, Takrama JS, Bernaet H, De Vuyst L. 2008b. Fermentation of
468 cocoa beans: influence of microbial activities and polyphenol concentrations on the flavour of
469 chocolate. *J Sci Food Agric.* 88:2288–2297. doi:10.1002/jsfa.

470 Codex Alimentarius Commission. 2018. Report of the 12th session of the codex committee on
471 contaminants in foods. Rome, Italy.

472 Engbersen N, Gramlich A, Lopez M, Schwarz G, Hattendorf B, Gutierrez O, Schulin R. 2019.
473 Cadmium accumulation and allocation in different cacao cultivars. *Sci Total Environ.* 678:660–
474 670. doi:10.1016/j.scitotenv.2019.05.001.

475 European Commission. 2014. Commission Regulation (EU) No 488/2014 of 12 May 2014
476 amending Regulation (EC) No 1881/2006 as regards maximum levels of cadmium in foodstuffs.
477 *Off J Eur Union.* 138:75–79.

478 Jespersen L, Nielsen DS, Hønholt S, Jakobsen M. 2005. Occurrence and diversity of yeasts
479 involved in fermentation of West African cocoa beans. *FEMS Yeast Res.* 5(4–5):441–453.
480 doi:10.1016/j.femsyr.2004.11.002.

481 Lewis C, Lennon AM, Eudoxie G, Umaharan P. 2018. Genetic variation in bioaccumulation and
482 partitioning of cadmium in *Theobroma cacao* L. *Sci Total Environ.* 640–641:696–703.
483 doi:10.1016/j.scitotenv.2018.05.365.

484 Lima LJR, Almeida MH, Rob Nout MJ, Zwietering MH. 2011. *Theobroma cacao* L., “the food
485 of the gods”: Quality determinants of commercial cocoa beans, with particular reference to the
486 impact of fermentation. *Crit Rev Food Sci Nutr.* 51(8):731–761.
487 doi:10.1080/10408391003799913.

488 Meter A, Atkinson RJ, Laliberte B. 2019. Cadmium in cacao from Latin America and the
489 Caribbean - A review of research and Potential Mitigation Solutions. Rome: Bioversity
490 International.

491 Papalexandratou Z, Vrancken G, de Bruyne K, Vandamme P, de Vuyst L. 2011. Spontaneous
492 organic cocoa bean box fermentations in Brazil are characterized by a restricted species diversity

493 of lactic acid bacteria and acetic acid bacteria. *Food Microbiol.* 28(7):1326–1338.
494 doi:10.1016/j.fm.2011.06.003.

495 Ramtahal G, Yen IC, Bekele I, Bekele F, Wilson L, Maharaj K, Harrynanan L. 2016.
496 Relationships between cadmium in tissues of cacao trees and soils in plantations of Trinidad and
497 Tobago. *Food Sci. Nutr.* 7:37–43. doi:10.4236/fns.2016.71005.

498 Ramtahal G, Umaharan P, Hanuman A, Davis C, Ali L. 2019. The effectiveness of soil
499 amendments, biochar and lime, in mitigating cadmium bioaccumulation in *Theobroma cacao* L.
500 *Sci Total Environ.* 693:133563. doi:10.1016/j.scitotenv.2019.07.369.

501 Santner J, Larsen M, Kreuzeder A, Glud RN. 2015. Two decades of chemical imaging of solutes
502 in sediments and soils - a review. *Anal Chim Acta.* 878:9–42. doi:10.1016/j.aca.2015.02.006.

503 Schwan RF, Wheals AE. 2004. The microbiology of cocoa fermentation and its role in chocolate
504 quality. *Crit Rev Food Sci Nutr.* 44(4):205–221. doi:10.1080/10408690490464104.

505 Vanderschueren R, Argüello D, Blommaert H, Montalvo D, Barraza F, Maurice L, Schreck E,
506 Schulin R, Lewis C, Vazquez JL, Umaharan P, Chavez E, Sarret G, Smolders E. 2021.
507 Mitigating the level of cadmium in cacao products: Reviewing the transfer of cadmium from soil
508 to chocolate bar. *Sci Total Environ.* 781:146779. doi:10.1016/j.scitotenv.2021.146779.

509 Vanderschueren R, De Mesmaeker V, Mounicou S, Isaure M-P, Doelsch E, Montalvo D,
510 Delcour JA, Chavez E, Smolders E. 2020. The impact of fermentation on the distribution of
511 cadmium in cacao beans. *Food Res Int.* 127:108743. doi:10.1016/j.foodres.2019.108743.

512 Vanderschueren R, Montalvo D, De Ketelaere B, Delcour JA, Smolders E. 2019. The elemental

513 composition of chocolates is related to cacao content and origin: A multi-element fingerprinting
514 analysis of single origin chocolates. *J Food Compos Anal.* 83. doi:10.1016/j.jfca.2019.103277.

515 De Vuyst L, Weckx S. 2016. The cocoa bean fermentation process: from ecosystem analysis to
516 starter culture development. *J Appl Microbiol.* 121(1):5–17. doi:10.1111/jam.13045.

517 Zhou C, van de Velde S, Baeyens W, Gao Y. 2018. Comparison of Chelex based resins in
518 diffusive gradients in thin-film for high resolution assessment of metals. *Talanta.* 186:397–405.
519 doi:10.1016/j.talanta.2018.04.085.

520